



THE UNIVERSITY *of* EDINBURGH

Edinburgh Research Explorer

Nitride tuning of lanthanide chromites

Citation for published version:

Black, AP, Johnston, HE, Oró-Solé, J, Bozzo, B, Ritter, C, Frontera, C, Attfield, JP & Fuertes, A 2016, 'Nitride tuning of lanthanide chromites', *Chemical Communications*, vol. 52, no. 23, pp. 4317-20.
<https://doi.org/10.1039/c6cc00744a>

Digital Object Identifier (DOI):

[10.1039/c6cc00744a](https://doi.org/10.1039/c6cc00744a)

Link:

[Link to publication record in Edinburgh Research Explorer](#)

Document Version:

Peer reviewed version

Published In:

Chemical Communications

General rights

Copyright for the publications made accessible via the Edinburgh Research Explorer is retained by the author(s) and / or other copyright owners and it is a condition of accessing these publications that users recognise and abide by the legal requirements associated with these rights.

Take down policy

The University of Edinburgh has made every reasonable effort to ensure that Edinburgh Research Explorer content complies with UK legislation. If you believe that the public display of this file breaches copyright please contact openaccess@ed.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.



Nitride tuning of lanthanide chromites

Ashley P. Black, Hannah Johnston, Judith Oró-Solé, Bernat Bozzo, Clemens Ritter, Carlos Frontera, J.Paul Attfield and Amparo Fuertes

Supplementary Information

Experimental Details

The oxynitrides $\text{RCrO}_{3-x}\text{N}_x$ ($\text{R} = \text{La}, \text{Pr}, \text{Nd}$) were prepared by treatment under $\text{NH}_3(\text{g})$ flow of $600 \text{ cm}^3/\text{min}$ (Carbureros Metálicos, 99.9%) of RCrO_4 precursor oxides between 700 and 800°C , using several cycles of 10 h with intermediate regrinding. The RCrO_4 compounds were prepared using the Pechini method. La_2O_3 (Aldrich, 99.9 %), Pr_6O_{11} (Aldrich, 99.9 %) or Nd_2O_3 (Aldrich, 99.9 %) were first fired at 900°C during 12 hours and dissolved in HNO_3 0.1 M . Then $\text{Cr}(\text{NO}_3)_3 \cdot 9\text{H}_2\text{O}$ (Aldrich 99.99 %), $\text{C}_6\text{H}_8\text{O}_7$ (citric acid, Aldrich 99.5 %) and $\text{C}_2\text{H}_6\text{O}_2$ (ethylene glycol), were subsequently added to the solution in molar ratios $\text{R}:\text{Cr}:\text{C}_6\text{H}_8\text{O}_7:\text{C}_2\text{H}_6\text{O}_2=1:1:1:1$, with continuous stirring and heating at $55\text{--}80^\circ\text{C}$. The solutions were evaporated for 12 h to form the precursor resins that were subsequently treated in air at 540°C for 15 hours .

N contents were determined by combustion analysis in oxygen in a Thermo Fisher Scientific instrument, heating the samples in oxygen up to 1060°C and using MgO , WO_3 and Sn as additives and atropine as a reference standard.

X-ray powder diffraction data were collected on a Siemens D5000 diffractometer using $\text{Cu K}\alpha$ radiation ($\lambda = 1.5418 \text{ \AA}$). Synchrotron X-ray powder diffraction data at room temperature were measured from capillary (0.3 mm diameter) samples in the angular range $1.038^\circ \leq 2\theta \leq 61.09^\circ$ at the MSPD beamline¹ of the ALBA Synchrotron (Cerdanyola del Vallès, Spain). Using a double Si (111) crystal monochromator, a short wavelength was selected and calibrated with Si NIST ($\lambda = 0.619714 \text{ \AA}$). Rietveld analysis was carried out using the program Fullprof.² Background refinement was performed by linear interpolation and the data were corrected for absorption.

Room temperature neutron powder diffraction data for $\text{LaCrO}_{2.72}\text{N}_{0.28}$ was collected on the D2B diffractometer at the Institut Laue-Langevin (ILL), Grenoble using 450 mg of sample placed in a 5 mm diameter vanadium can. The neutron wavelength was 1.594 \AA . Room temperature data was collected for $\text{NdCrO}_{2.58}\text{N}_{0.42}$ and $\text{PrCrO}_{2.64}\text{N}_{0.36}$ on the D20 diffractometer also at the Institut Laue-Langevin. The neutron wavelength was 1.36 \AA using a take-off angle of 118° . Low temperature neutron powder diffraction was carried out on all the samples on D1B diffractometer at the ILL. The neutron wavelength was 2.524 \AA . For the $\text{R} = \text{La}$ and Nd samples a series of scans were taken at intervals of approximately 6.5 K between $10 \text{ K} - 315 \text{ K}$ for $\text{R} = \text{La}$ and approximately 3.5 K between $1.5 \text{ K} - 215 \text{ K}$ for $\text{R} = \text{Nd}$. For $\text{R} = \text{Pr}$, data was collected in 50 K intervals between 1.5 K and 250 K . Powder diffraction data were analysed using the FullProf software package.² The anion composition of the oxynitrides was constrained by the chemically determined value.

Magnetic measurements were performed between at $H=2000 \text{ G}$ between 10 K and 400 K using a Quantum Design SQUID magnetometer.

Table S1. Summary of the *Pbnm* model for LaCrO_{2.72}N_{0.28} refined against room temperature synchrotron X-ray powder diffraction data. Refined cell parameters: a=5.52201(4), b=5.48192(5), c=7.6573(7) Å. R_{wp} = 4.33 %, χ^2 = 8.21 for 59 variables.

Atom	Site	x	y	z	B _{iso} (Å ²)
La	4c	0.9978(2)	0.01274(9)	1/4	0.715(6)
Cr	4b	1/2	0.0	0.0	0.344(8)
X1	4b	0.070(1)	0.493(1)	1/4	0.31(4)
X2	8d	0.727(1)	0.268(1)	0.0353(5)	0.31(4)
		La-X1	La-X2	Cr-X1	Cr-X2
Bond length / Å		2.391(6)	2.474(5) × 2	1.980(1) × 2	1.953(6) × 2
		2.660(6)	2.640(6) × 2		1.989(6) × 2
		2.879(6)	2.821(5) × 2		
		3.136(6)	3.096(5) × 2		
		Cr-X1-Cr	Cr-X2-Cr		
Bond Angle / °		157.45(5)	161.5(3)		

Table S2. Summary of the *Pbnm* model for PrCrO_{2.81}N_{0.19} refined against room temperature synchrotron X-ray powder diffraction data. Refined cell parameters: a= 5.45535(4), b= 5.49076(3), c= 7.72751(5) Å. R_{wp} = 3.24 %, χ^2 = 3.93 for 55 variables.

Atom	Site	x	y	z	B _{iso} (Å ²)
Pr	4c	0.9932(1)	0.03577(4)	1/4	0.610(4)
Cr	4b	1/2	0.0	0.0	0.158(8)
X1	4b	0.0736(9)	0.4815(5)	1/4	0.25(6)
X2	8d	0.7147(7)	0.2939(6)	0.0388(5)	0.25(6)
		Pr-X1	Pr-X2	Cr-X1	Cr-X2
Bond length / Å		2.382(5)	2.390(4) × 2	1.976(1) × 2	1.948(4) × 2
		2.486(3)	2.642(4) × 2		2.016(3) × 2
		3.075(3)	2.705(4) × 2		
		3.107(5)	3.286(4) × 2		
		Cr-X1-Cr	Cr-X2-Cr		
Bond Angle / °		155.80(4)	155.0(2)		

Table S3. Summary of the *Pbnm* model for NdCrO_{2.58}N_{0.42} refined against room temperature synchrotron X-ray powder diffraction data. Refined cell parameters: a=5.42787(3), b=5.50102(3), c=7.70936(4) Å. R_{wp} = 2.36 %, χ^2 = 7.72 for 96 variables.

Atom	Site	x	y	z	B _{iso} (Å ²)
Nd	4c	0.9919(1)	0.04131(5)	1/4	0.787(7)
Cr	4b	1/2	0.0	0.0	0.17(1)
X1	4b	0.0851(9)	0.484(7)	1/4	0.53(4)
X2	8d	0.7190(7)	0.2935(7)	0.0410(5)	0.43(4)
		Nd-X1	Nd-X2	Cr-X1	Cr-X2
Bond length / Å		2.318(5)	2.401(4) × 2	1.984(1) × 2	1.928(4) × 2
		2.487(3)	2.591(4) × 2		2.030(4) × 2
		3.108(3)	2.716 (4) × 2		
		3.148(5)	3.300 (4) × 2		
		Cr-X1-Cr	Cr-X2-Cr		
Bond Angle / °		152.57(5)	155.0(2)		

Table S4. Summary of the *Pbnm* model for LaCrO_{2.72}N_{0.28} refined against room temperature neutron powder diffraction data. Refined cell parameters: a=5.5237(2), b=5.4787(2), c=7.7684(2) Å. R_{wp} = 2.45 %, χ^2 = 1.89 for 41 variables.

Atom	Site	x	y	z	B _{iso} (Å ²)
La	4c	0.0037(8)	0.0152(7)	1/4	0.038(4)
Cr	4b	1/2	0.0	0.0	0.036(5)
X1	4b	0.0671(7)	0.487(1)	1/4	0.732(4)
X2	8d	0.7325(7)	0.2705(7)	0.0362(3)	0.732(4)
		La-X1	La-X2	Cr-X1	Cr-X2
Bond length / Å		2.376(6)	2.502(4) × 2	1.978(8) × 2	1.961(4) × 2
		2.613(8)	2.638(5) × 2		1.981(4) × 2
		2.911(8)	2.814(4) × 2		
		3.156(6)	3.085(4) × 2		
		Cr-X1-Cr	Cr-X2-Cr		
Bond Angle / °		158.05(3)	161.5(2)		

Table S5. Summary of the *Pbnm* model for $\text{PrCrO}_{2.64}\text{N}_{0.36}$ refined against room temperature neutron powder diffraction data. Refined cell parameters: $a = 5.3976(2)$, $b = 5.4336(2)$, $c = 7.6461(3)$ Å. $R_{wp} = 1.40\%$, $\chi^2 = 1.82$ for 128 variables.

Atom	Site	x	y	z	B_{iso} (Å ²)
Pr	4c	0.9959(2)	0.0302(8)	1/4	0.341(7)
Cr	4b	1/2	0.0	0.0	0.599(4)
X1	4b	0.0756(9)	0.4843(7)	1/4	0.599(4)
X2	8d	0.7106(5)	0.2904(5)	0.0403(4)	0.599(4)
		Pr-X1	Pr-X2	Cr-X1	Cr-X2
Bond length / Å		2.326(1)	$2.348(6) \times 2$	$1.956(1) \times 2$	$1.958(3) \times 2$
		2.504(6)	$2.635(7) \times 2$		$1.969(3) \times 2$
		2.997(6)	$2.687(5) \times 2$		
		3.095(1)	$3.236(6) \times 2$		
		Cr-X1-Cr	Cr-X2-Cr		
Bond Angle / °		155.4(4)	154.5(1)		

Table S6. Summary of the *Pbnm* model for $\text{NdCrO}_{2.58}\text{N}_{0.42}$ refined against room temperature neutron powder diffraction data. Refined cell parameters: $a = 5.3715(2)$, $b = 5.4459(2)$, $c = 7.6316(2)$ Å. $R_{wp} = 1.36\%$, $\chi^2 = 2.07$ for 93 variables.

Atom	Site	x	y	z	B_{iso} (Å ²)
Nd	4c	0.9930(9)	0.0388(5)	1/4	0.182(3)
Cr	4b	1/2	0.0	0.0	0.182(3)
X1	4b	0.0796(8)	0.4800(7)	1/4	0.897(4)
X2	8d	0.7097(5)	0.2944(5)	0.0421(4)	0.897(4)
		Nd-X1	Nd-X2	Cr-X1	Cr-X2
Bond length / Å		2.319(6)	$2.340(4) \times 2$	$1.958(1) \times 2$	$1.946(3) \times 2$
		2.447(5)	$2.602(4) \times 2$		$1.985(3) \times 2$
		3.078(5)	$2.674(4) \times 2$		
		3.092(6)	$3.288(4) \times 2$		
		Cr-X1-Cr	Cr-X2-Cr		
Bond Angle / °		153.99(4)	153.16(1)		

Table S7. Summary of the magnetic refinements for *Pbnm* models at base temperatures using D1B neutron diffraction data. Refined lattice parameters and ordered moments at Cr and Nd sites are shown.

Sample	T/K	a/Å	b/Å	c/Å	$\mu_y(\text{Cr})/\mu_B$	$\mu_z(\text{Ln})/\mu_B$
LaCrO _{2.72} N _{0.28}	10	5.4508(9)	5.4086(6)	7.666(2)	2.73(5)	
PrCrO _{2.64} N _{0.36}	1.5	5.3258(7)	5.4039(7)	7.575(1)	2.60(8)	
NdCrO _{2.58} N _{0.42}	1.5	5.3569(7)	5.3922(6)	7.598(1)	2.59(5)	1.4(1)

Table S8 Values of Curie temperatures and other magnetic parameters from Curie-Weiss fits to data above T_C for Ln = Pr and Nd samples (this was not possible for Ln = La due to the high T_C). Effective Cr paramagnetic moments were estimated by subtracting the ideal Ln^{3+} contribution from the total as shown in the Table footnote.

Ln, x ^[a]	T_C/K	T_{SR}/K	θ/K	μ_{eff}/μ_B	$\mu_{\text{eff}}(\text{Cr})/\mu_B^{[a]}$
La, 0	293				
La, 0.11	285				
La, 0.17	283				
La, 0.21	283				
La, 0.25	280				
La, 0.28	281				
Pr, 0	240	-	-217	5.50	4.17
Pr, 0.19	229	-	-156	5.17	3.73
Pr, 0.36	232	-	-43	4.30	2.38
Nd, 0	226	28	-251	4.72	3.03
Nd, 0.35	210	44	-138	5.21	3.75
Nd, 0.59	214	-	-91	4.60	2.84

^[a] $\mu_{\text{eff}}(\text{Cr}) = [\mu_{\text{eff}}^2 - \mu_{\text{eff}}(\text{Ln}^{3+})^2]^{1/2}$ where $\mu_{\text{eff}}(\text{Pr}^{3+}) = 3.58$ and $\mu_{\text{eff}}(\text{Nd}^{3+}) = 3.62 \mu_B$.




Figure S1. Observed and calculated synchrotron X-ray powder diffraction patterns for $\text{PrCrO}_{2.81}\text{N}_{0.19}$.




Figure S2. Observed and calculated synchrotron X-ray powder diffraction patterns for $\text{NdCrO}_{2.58}\text{N}_{0.42}$.




Figure S3. Observed and calculated 10 K neutron powder diffraction patterns for $\text{LaCrO}_{2.72}\text{N}_{0.28}$ ($\lambda=2.524 \text{ \AA}$).




Figure S4. Observed and calculated 1.5 K neutron powder diffraction patterns for $\text{PrCrO}_{2.64}\text{N}_{0.36}$ ($\lambda=2.524 \text{ \AA}$).




Figure S5. Observed and calculated room temperature neutron powder diffraction patterns for $\text{PrCrO}_{2.64}\text{N}_{0.36}$ ($\lambda = 1.36 \text{ \AA}$)




Figure S6. Observed and calculated room temperature neutron powder diffraction patterns for $\text{NdCrO}_{2.58}\text{N}_{0.42}$ ($\lambda = 1.36 \text{ \AA}$).




Figure S7. Thermal variation of refined Cr magnetic moment for $\text{LaCrO}_{2.72}\text{N}_{0.28}$.




Figure S8. Thermal variation of refined Cr magnetic moment for $\text{PrCrO}_{2.64}\text{N}_{0.36}$.

¹ F. Fauth, I. Peral, C. Popescu, C. and M. Knapp, M. *Powder Diffraction* **28**, S360–S370 (2013).

² J. Rodríguez-Carvajal, *Phys. B*, 1993, **192**, 55.